

Use of a Treated, Devitalized and Sterile Umbilical Cord Amniotic Membrane in the Treatment of PIP Joint Osteoarthritis

NCT06034262

Status	RECRUITING
Phase	Phase 2
Sponsor	TBF Genie Tissulaire
Enrollment	14 participants

Key Eligibility Criteria

Inclusion (7)

- Male or female; age between 18 and 75 years.
- Patient with symptomatic PIPJ OA.
- Patient treated medically for functional signs for more than 3 months, without improvement, justifying a surgical intervention.
- Patient with a QuickDash score \geq 33 points (converted to 50%).
- Patient with pain, at rest, in the PIPJ \geq 4/10 on a VAS.

... and 2 more (see full listing online)

Exclusion (5)

- Pregnant or breastfeeding woman; woman without effective contraception.
- Patient with signs of neuropathy with functional disorders such as hyperesthesia.
- Patient with uncontrolled epilepsy or neuropathy that contraindicates locoregional anesthesia.
- Person confined by a judicial or administrative decision.
- Adult subjected to legal protection measures or unable to provide his/her consent.

Locations (3 total)

CHU de Besançon, Besançon, France
Institut Chirurgical de la Main et du Membre Supérieur, Lyon, France
Centre de la Main - Clinique Jules Verne, Nantes, France