

# Use of the Neurolyser XR for the Treatment of Low Back Pain Related to Sacroiliitis

NCT06053255

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	FUSMobile Inc.
Enrollment	10 participants

## Key Eligibility Criteria

---

### Inclusion (12)

- Men and women aged 18 and older
- BPI  $\leq$  40
- Patients who are able and willing to give consent and able to attend all study visits.
- Patients who are able to communicate with the treating physician.
- Patients must have chronic lower back pain (LBP) attributed to degenerative sacroiliitis diagnosed by a pain physician using the following guidelines:  
... and 7 more (see full listing online)

### Exclusion (29)

- Patients on dialysis
- Patients with evidence of lumbosacral radiculopathy on MRI, CT or physical exam findings, including radicular leg pain (symptomatic moderate or severe foraminal or central canal stenosis).
- Spinal pathology that may impede recovery such as spina bifida occulta, grade II or higher.
- Spondylolisthesis at L5/S1, or scoliosis
- Patients with radiofrequency ablation (RF/RFA) treatment for sacroiliitis LBP  
... and 24 more (see full listing online)

## Locations (1 total)

---

University of Virginia, Department of Anesthesiology, Charlottesville, Virginia, United States