

Two Electroacupuncture Waveforms for Different Severity Groups of Bell Palsy

NCT06063954

Status	RECRUITING
Phase	Not Applicable
Sponsor	The Third Affiliated hospital of Zhejiang Chinese Medical University
Enrollment	120 participants

Key Eligibility Criteria

Inclusion (5)

- Diagnosed as BP by specialist.
- The score of FNGS 2.0 e 15 at the day 21 since the onset of BP.
- years d age d 65 years.
- Received prednisolone within 72 hours since initial symptoms of BP, the prednisolone dose used was 60 mg per day for 5 days and then reduced by 10 mg per day.
- Signed informed consent and volunteered to participate in this study.

Exclusion (9)

- Facial palsy caused by other diseases or injury.
 - Ramsey-Hunt syndrome.
 - Bilateral facial palsy.
 - History of previous facial palsy.
 - Manifesting facial spasm, facial synkinesis or contracture at day 21 since the onset of BP.
- ... and 4 more (see full listing online)

Locations (3 total)

The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China
The Third Affiliated hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China
Hangzhou First People' s Hospital, Hangzhou, Hangzhou, Zhejiang, China