

Timing of CHolecystectomy In Severe PANcreatitis

NCT06113419

Status	RECRUITING
Phase	Not Applicable
Sponsor	Hospital Universitario Mayor Méderi
Enrollment	134 participants

Key Eligibility Criteria

Inclusion (3)

- Age ≥18 years, diagnosis of pancreatitis according to Atlanta guidelines, moderately severe or severe pancreatitis (APACHE score ≥8 on admittance)
- Biliary pancreatitis diagnosed on imaging (be it ultrasound, magnetic resonance imaging and/or tomography)
- Recovery of pancreatitis by tolerance of oral intake (defined as 24 hours of food consumption of any consistency without emetic episodes and pain defined as 4/10 on the visual analogue score of pain) and written informed consent.

Exclusion (9)

- Pregnancy
- History of cholecystectomy
- Planned open cholecystectomy
- Pancreatitis-associated complication before laparoscopic cholecystectomy (compartment syndrome, bleeding and/or need for peripancreatic collection drainage)
- Chronic pancreatitis,
- ... and 4 more (see full listing online)

Locations (2 total)

Hospital Universitario Mayor-Méderi, Bogota, Cundinamarca, Colombia
Hospital Universitario Méderi Barrios Unidos, Bogota, Cundinamarca, Colombia