

Study of the Telitaccept in Pediatric Patients With Frequently Relapsing or Steroid Dependent Nephrotic Syndrome

NCT06125405

Status	RECRUITING
Phase	Phase 3
Sponsor	The Children's Hospital of Zhejiang University School of Medicine
Enrollment	20 participants

Key Eligibility Criteria

Inclusion (5)

- Sensitive but frequent relapses or steroids dependence nephrotic syndrome
- Age: 2 to 18 years old
- Normal renal function: estimated glomerular filtration rate ≥ 90 ml/min/1.73m²
- Morning urine protein $\leq 1+$ or urine protein-creatinine ratio ≤ 0.2 g/g (≤ 20 mg/mmol) for 3 consecutive days and above when in enroll
- No rituximab was used within 6 months, no tacrolimus, mycophenolate mofetil, cyclosporine A, or cyclophosphamide was used within 3 months, no ACTH was used within 3 months prior to the enrollment

Exclusion (11)

- Family history of nephrotic syndrome, chronic glomerulonephritis or uremia
- Leukopenia (White Blood Cells $\leq 3.0 \times 10^9$ / L)
- Moderate to severe anemia (hemoglobin ≤ 9.0 g/dL)
- Thrombocytopenia (platelet count $\leq 100 \times 10^12$ / L)
- Positive Hepatitis B virus serological indicators (Hepatitis B surface antigen or / and Hepatitis B virus e antigen or / and Hepatitis B core antibody), Hepatitis C virus-positive or patients with abnormal liver function (2 or more times of alanine aminotransferase or total bilirubin was exceeded the normal value, and continued to rise for 2 weeks)

... and 6 more (see full listing online)

Locations (1 total)

Children's Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China