

Assessing Interventions to Increase Tdap Acceptance for Non-birthing Partners in Pregnancy

NCT06135636

Status	RECRUITING
Phase	Phase 4
Sponsor	Women and Infants Hospital of Rhode Island
Enrollment	150 participants

Key Eligibility Criteria

Inclusion (4)

- Partners of pregnant patients who are receiving their prenatal care at the OGCC
- years old: This age range was selected as 19 years is both the age at which adult Tdap vaccination is recommended by the CDC as well as the age for the State supplied adult Tdap and Influenza Vaccines. Fifty years old is the upper limit of "reproductively aged individuals, the target population for this study.
- Have not or are unsure if they have received the adult Tdap vaccine or booster in the last 10 years: In addition to the CDC recommendation for adult Tdap vaccination and Td or Tdap booster every 10 years, the CDC recommends vaccination for any adult who is unsure of their vaccination status to ensure they are vaccinated as repeat vaccination does not cause additional harm.
- Fluency in English or Spanish: Consents, surveys, and Tdap and cocooning information will be available in both languages.

Exclusion (2)

- Latex allergy: Contraindication to the state supplied Tdap vaccine
- Lethal fetal anomaly diagnosed prior to enrollment to prevent undue distress with follow-up postpartum

Locations (1 total)

Women and Infants Hospital, Providence, Rhode Island, United States