

# Exploring the Benefit of Peripheral Nerve Stimulation in Treating Pain From Chemo-induced Peripheral Neuropathy: A Longitudinal Single Center Feasibility Study

NCT06162403

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	M.D. Anderson Cancer Center
<b>Enrollment</b>	10 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- Participants diagnosed with chronic (≥90 days duration) CIPN (due to either vinca alkaloids, taxanes, bortezomib, thalidomide, platinum-based compounds or ionizing irradiation) of the lower extremity, seen at Pain Management Center at MD Anderson Cancer Center
- Participants reports baseline pain ≥ 4 (0-10 scale, NRS)
- Participants between ages 18-85 years old
- Participants who have completed chemotherapy within the previous year at the time of enrollment

### Exclusion (5)

- Participants with cognitive dysfunction
- Participants with recent history (<6 months) of drug or alcohol abuse
- Participants with open skin lesion or undergoing antibiotic therapy for local or systemic infection
- Participants with allergies to local anesthesia, steroids, or adhesives
- Participants with conditions that conflict with the SPRINT PNS System Indications for Use, including Contraindications and Warnings.

## Locations (1 total)

---

MD Anderson Cancer Center, Houston, Texas, United States