

The Effects of Remimazolam on the Incidence of Severe Hypoxia During Sedated Hysteroscopy for Assisted Reproduction in Overweight or Obese Patients

NCT06187896

Status	RECRUITING
Phase	Not Applicable
Sponsor	RenJi Hospital
Enrollment	600 participants

Key Eligibility Criteria

Inclusion (5)

- Age between 20 and 50 years.
- Patients undergoing sedated hysteroscopy for assisted reproduction.
- ASA classification I-II.
- BMI \leq 23 kg/m².
- Patients who have signed an informed consent form.

Exclusion (10)

- SpO₂ $<$ 95% in patients inhaling air upon entering the room.
- Diagnosed with chronic obstructive pulmonary disease (COPD) or currently suffering from other acute and chronic lung diseases requiring long-term or intermittent oxygen therapy.
- Patients with a history of mental and neurological disorders: such as depression, severe central nervous system depression, Parkinson's disease, basal ganglia disease, schizophrenia, epilepsy, Alzheimer's disease, myasthenia gravis, etc.
- Patients with severe liver dysfunction.
- Patients with severe renal insufficiency (requiring dialysis before surgery).

... and 5 more (see full listing online)

Locations (1 total)

Diansan Su, Shanghai, Shanghai Municipality, China