

# Usefulness of Corifollitropin as Alternative to Conventional Daily rFSH Protocols in Oocyte Donors Undergoing Pituitary Suppression With Medroxyprogesterona Acetate (MPA)

NCT06193135

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Instituto Valenciano de Infertilidad, IVI VALENCIA
<b>Enrollment</b>	318 participants

## Key Eligibility Criteria

---

### Inclusion (12)

- Signature of the subject's informed consent prior to any trial-related activity.
  - Age between 18 and 35 years (both inclusive).
  - Regular menstrual cycle, from 25 to 35 days (both inclusive).
  - Absence of physical and psychological illness at the time of donation at the discretion of the investigator.
  - BMI 18-28 kg/m2 (both inclusive) at the time of donation.
- ... and 7 more (see full listing online)

### Exclusion (9)

- Concurrent participation in another clinical trial.
  - Previous participation in this clinical trial.
  - Use of long-term hormonal contraception (hormonal IUD or subcutaneous implants) at least 1 month prior to enrollment.
  - Any systemic or metabolic disorder (i.e.: diabetes...) that contraindicates the use of gonadotropins.
  - Personal history of thrombophlebitis and thromboembolic phenomena and hypertension.
- ... and 4 more (see full listing online)

## Locations (2 total)

---

IVI Valencia, Valencia, Valencia, Spain  
Ivi Valencia, Valencia, Valencia, Spain