

# RIVaroxaban Versus Low-molecular Weight Heparin in Patients With Lower Limb Trauma Requiring Brace or CASTing

NCT06195540

---

Status	RECRUITING
Phase	Phase 3
Sponsor	University Hospital, Angers
Enrollment	1,424 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Patient aged 18 or over ;
- Consultation in an emergency department of a participating centre;
- Trauma to the lower limb requiring rigid or semi-rigid orthopaedic immobilisation;
- Expected duration of orthopaedic immobilisation of at least 2 weeks;
- TRiP(cast) score  $\geq 7$  ;

... and 2 more (see full listing online)

### Exclusion (6)

- Patient that have to be hospitalized after emergency department for other reason than lower limb trauma
- Active bleeding or high risk of bleeding,
- Known contraindication to rivaroxaban or LMWH;
- Taking any anticoagulant or antiplatelet agent before the trauma (only antithrombotic authorised: aspirin  $< 325\text{mg/d}$ );
- Pregnant or breastfeeding woman;

... and 1 more (see full listing online)

## Locations (35 total)

---

Agen-Nerac Hospital, Emergency Department, Agen, France  
Angers University Hospital, Emergency department, Angers, France  
Argenteuil hospital, Emergency department, Argenteuil, France

... and 32 more locations