

The Role of Flavor in the Substitutability of E-cigarettes for Combustible Cigarettes Among Persistent Smokers

NCT06264154

Status	RECRUITING
Phase	Not Applicable
Sponsor	Abramson Cancer Center at Penn Medicine
Enrollment	210 participants

Key Eligibility Criteria

Inclusion (9)

- Able to communicate fluently in English (i.e. speaking, writing, and reading)
- Male and female smokers who are \geq 21 years of age and self-report smoking at least 5 cigarettes (menthol and/or non-menthol) per day for at least the last 12 months
- or more failed quit attempts and the use of smoking cessation medication on at least one prior attempt.
- Ever use of an e-cigarette
- Have a carbon monoxide (CO) greater than 10 ppm
- ... and 4 more (see full listing online)

Exclusion (20)

- Smoking Behavior Regular use of nicotine-containing products other than cigarettes (e.g., chewing tobacco, snuff, snus, cigars, e-cigs, etc.). Regular e-cigarette use is defined as greater than 5 days/past 30 days
- Participants agreeing to abstain from using nicotine-containing products other than cigarettes for the duration of the study will be considered eligible.
- Current enrollment or plans to enroll in a smoking cessation program over the duration of the study.
- Current use of smoking cessation medication
- Provide a CO breath test reading less than 10 ppm at Intake.
- ... and 15 more (see full listing online)

Locations (1 total)

University of Pennsylvania, Philadelphia, Pennsylvania, United States