

Particulate vs. Non-Particulate Steroid for Sacroiliac Joint Injection

NCT06268704

Status	RECRUITING
Phase	Phase 4
Sponsor	University of New Mexico
Enrollment	230 participants

Key Eligibility Criteria

Inclusion (6)

- Aged ≥ 18 , capable of understanding and providing consent in English, capable of complying with the outcome instruments used, capable of attending all planned follow up visits
- Unilateral low back/buttocks pain of at least 2 weeks.
- Patient reported 7 day average of numeric pain rating score (NPRS) low back/buttocks pain of at least 5/10 at baseline evaluation
- Clinical diagnosis of sacroiliac joint pain as diagnosed by a board certified Physiatrist including history of low back/buttocks pain and at least 2 positive physical exam findings (including positive fortin finger sign, pain with palpation of posterior superior iliac spine, positive FABER's test, positive Gaenslan's test, positive sacral distraction, positive thigh thrust, positive lateral compression, positive sacral thrust)
- Patient consents to sacroiliac joint corticosteroid injection in a shared decision-making process with the treating physician.

... and 1 more (see full listing online)

Exclusion (21)

- Clinical suspicion of alternative process is greater than clinical suspicion of sacroiliac joint pain
- Those receiving remuneration for their pain treatment (e.g., disability, worker's compensation).
- Those involved in active litigation relevant to their pain.
- Those unable to read English and complete the assessment instruments.
- Those unable to attend follow up appointments

... and 16 more (see full listing online)

Locations (1 total)

University of New Mexico Hospital, Albuquerque, New Mexico, United States