

Efficacy of Intraoperative Use of 20% Albumin Combined With Ringer Lactate Versus Ringer Lactate During Cytoreductive Surgery With Hyperthermic Intraperitoneal Chemotherapy

NCT06351475

Status	RECRUITING
Phase	Not Applicable
Sponsor	Assistance Publique - Hôpitaux de Paris
Enrollment	140 participants

Key Eligibility Criteria

Inclusion (4)

- Male and female patients aged 18 years old and older
- Planned cytoreductive surgery with HIPEC for peritoneal carcinomatosis (from gynaecological or digestive origin)
- Patient volunteer for the study and provided written informed consent
- Patient affiliated to the French Health Insurance

Exclusion (19)

- Adults ≥ 75 years old
- Functional status rendering the patient ineligible for cytoreduction with CHIP (ECOG ≥ 2 or Karnofsky index < 75)
- Pre-existing preoperative conditions affecting albuminemia (hepatic cirrhosis, nephrotic syndrome, exudative enteropathy, malnutrition)
- Severe preoperative hypoalbuminemia (albuminemia < 20 g/L)
- History of chronic left heart failure with reduced left ventricular ejection fraction (left ventricular ejection fraction $< 40\%$)
- ... and 14 more (see full listing online)

Locations (7 total)

Institut régional du Cancer de Montpellier, Montpellier, France
La Pitié Salpêtrière hospital, Paris, France
Lariboisière hospital, Paris, France
... and 4 more locations