

Efficacy, Safety, and Acceptability of Mifepristone 50 mg Once-weekly as a Contraceptive

NCT06394999

Status	RECRUITING
Phase	Phase 3
Sponsor	Leiden University Medical Center
Enrollment	1,186 participants

Key Eligibility Criteria

Inclusion (13)

- Age 18-35 at the moment of signing the informed consent.
- In case of 30 years or older, prior HPV or pap-test.
- Understand and speak Dutch or English.
- Willing to use mifepristone as the only method of contraception for 12 months.
- Able to take oral medication and willing to adhere to the study protocol.

... and 8 more (see full listing online)

Exclusion (17)

- Currently pregnant or breast-feeding.
- Desire to become pregnant within the following 12 months.
- Signs of current endometritis, incomplete abortion after miscarriage or induced abortion, or retained pla-cental rests after delivery.
- Undiagnosed post-coital bleeding or unscheduled bleeding (spotting with COC, patch, vaginal ring, hormo-nal IUD, implant allowed).
- Known subfertility or history of ectopic pregnancy, unless intra-uterine pregnancy afterwards, or grade 3-4 endometriosis proven by laparoscopy.

... and 12 more (see full listing online)

Locations (14 total)

Noordwest Ziekenhuisgroep, Alkmaar, Netherlands
Ziekenhuisgroep Twente, Almelo, Netherlands
Flevoziekenhuis, Almere Stad, Netherlands
... and 11 more locations