

FLudrocortisone Administration in Aneurysmal Subarachnoid Haemorrhage

NCT06409364

Status	RECRUITING
Phase	Phase 2
Sponsor	The George Institute
Enrollment	524 participants

Key Eligibility Criteria

Inclusion (5)

- Age 18 years or older
- Diagnosed with subarachnoid haemorrhage from an aneurysm confirmed on computed tomography angiography (CTA) or digital subtraction angiography (DSA) of the intra-cranial arteries
- Aneurysm has been secured
- Hospital admission for aSAH within 96 hours
- Currently being treated in a critical care environment

Exclusion (8)

- Unable to receive enteral medications
 - Pre-existing glucocorticoid or mineralocorticoid treatment
 - Previous allergic reaction to fludrocortisone
 - History of cardiac, hepatic, or renal failure
 - Hyponatremia or hyponatremia ($Na^+ > 145 \text{mmol/L}$ or $Na^+ < 125 \text{mmol/L}$) on the most recent blood sample at the time of screening.
- ... and 3 more (see full listing online)

Locations (16 total)

John Hunter Hospital, New Lambton Heights, New South Wales, Australia
Nepean Hospital, Penrith, New South Wales, Australia
Prince of Wales Hospital, Randwick, New South Wales, Australia
... and 13 more locations