

# Exercise Training Study of Patients With Claudicatio Intermittens

NCT06448390

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Norwegian University of Science and Technology
<b>Enrollment</b>	100 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Pain in the calves after 300 m, or less measured on a treadmill
  - Stable symptoms last 6 months or more
  - The pain must be gone within 5 minutes after the patients stop walking
  - Ankle-Brachial index at 0.7 or lower at rest
  - Using statins and antiplatelet drugs since at least 3 months
- ... and 2 more (see full listing online)

### Exclusion (13)

- pain in the hips or elsewhere outside the calves while walking
  - Remaining pain more than 5 minutes after stopped walking.
  - Impossible to compress the arteries while measuring ABI or ABI lower than 0.4
  - Use of statins and antiplatelet drugs less than 3 months
  - symptoms less than 6 months
- ... and 8 more (see full listing online)

## Locations (14 total)

---

Ålesund Sykehus, Ålesund, Norway  
Haukeland University Hospital, Bergen, Norway  
Vestre Viken, Drammen sykehus, Drammen, Norway  
... and 11 more locations