

Intra-articular Allocetra in Osteoarthritis of the 1st Carpo-metacarpal Joint

NCT06459063

Status	RECRUITING
Phase	Phase 1, Phase 2
Sponsor	Kaplan Medical Center
Enrollment	46 participants

Key Eligibility Criteria

Inclusion (6)

- Subjects eligible for this clinical study must fulfill all of the following:
 - Age 40 years or older.
 - Patients with OA of the first CMC joint of the target thumb (basal thumb joint) who have failed conventional therapies, with pain assessed when not taking analgesic medications.
 - Score of 6 or higher on the Functional Index for Hand Osteoarthritis (FIHOA).
 - X-ray confirming OA of the first CMC joint of the target thumb with a Grade of 2 or 3 according to Eaton classification.
- ... and 1 more (see full listing online)

Exclusion (9)

- Subjects not eligible for this study include those that have any of the following:
 - Any significant injury, fracture, surgery, active local infection, deformity, severe Carpal Tunnel Syndrome (CTS), DeQuervain's tenosynovitis, trigger finger, or a ganglion cyst of the target hand.
 - History of chondrocalcinosis in the target joint, concomitant rheumatic disease.
 - Previous intra-articular injection of steroid, hyaluronate, or other agent, into the target joint within 3 months prior to screening visit.
 - Other limb pain of unknown etiology, or clinically significant widespread pain syndrome, e.g., fibromyalgia.
- ... and 4 more (see full listing online)

Locations (1 total)

Kaplan Medical Center, Rehovot, Israel