

# Application of Autologous Adipose Stromal Vascular Fraction in the Treatment of Premature Ovarian Insufficiency

NCT06481969

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Rui Yang
Enrollment	260 participants

## Key Eligibility Criteria

---

### Inclusion (6)

- Women aged e20 and d39 years who have childbearing requirements.
  - Cessation of menstruation or oligomenorrhea for at least 4 months
  - Serum level of basal follicle stimulating hormone (FSH) \>25 U/L at least 2 times (during the second to fourth day of the menstrual cycle or during amenorrhea; The interval between the two tests is 4 weeks)
  - Serum level of Anti-Mullerian hormone (AMH) d1.1 ng/ml
  - Women with intact uterus and bilateral adnexa.
- ... and 1 more (see full listing online)

### Exclusion (9)

- Women with autoimmune diseases.
  - Women with abnormal and uncontrolled thyroid function.
  - Women with tumors in bilateral adnexa that are not clearly benign or malignant.
  - Women with a history of malignant tumors, radiation therapy or chemotherapy.
  - Women with a history of venous thrombosis or pulmonary embolism during the screening period.
- ... and 4 more (see full listing online)

## Locations (1 total)

---

Peking University Third Hospital, Beijing, Beijing Municipality, China