

Effects of Vagus Nerve Pulmonary Branch Block on Postoperative Cough After VATS Lung Resection

NCT06500949

Status	RECRUITING
Phase	Not Applicable
Sponsor	Second Affiliated Hospital, School of Medicine, Zhejiang University
Enrollment	104 participants

Key Eligibility Criteria

Inclusion (7)

- Age between 18 and 70 years old.
- BMI between 18 and 30 kilograms per square meter (kg/m²).
- ASA Physical Status Classification of I, II, or III.
- Preoperative pulmonary imaging demonstrating peripheral lesions, with clinical staging Td2, Nd1, M0.
- Undergone thoracoscopic lung resection surgery.

... and 2 more (see full listing online)

Exclusion (5)

- Patients with a history of chemotherapy or previous pulmonary surgery.
- Presence of chronic cough due to respiratory infectious diseases, pharyngitis, rhinitis, COPD, asthma, post-nasal drip syndrome, etc.
- Individuals exhibiting ECG abnormalities such as atrial fibrillation, bundle branch block, frequent ventricular premature beats, pre-excitation syndrome, etc.
- Currently using ACE inhibitor medications.
- Presence of preoperative hoarseness.

Locations (1 total)

Second affiliated Hospital School of Medicine, Zhejiang University, Hangzhou, Zhejiang, China