

Effect of a GnRH Analog on Hepatic Steatosis

NCT06523530

Status	RECRUITING
Phase	Phase 4
Sponsor	Aristotle University Of Thessaloniki
Enrollment	62 participants

Key Eligibility Criteria

Inclusion (4)

- women of reproductive age
- diagnosis of endometriosis. The disease is suspected by patient's individual history (chronic pelvic pain, dyspareunia or/and dysmenorrhea) and the ultrasonographic imaging (chocolate cysts). The diagnosis is confirmed histologically, after laparoscopic surgical treatment and biopsy sampling, which will be interpreted by an independent blinded pathologist.
- use of contraceptives, which is the first line treatment, is contraindicated or the patient does not consent to receive contraceptives, due to personal preferences.
- written informed consent to participate to the study

Exclusion (10)

- mean ethanol consumption $\gt 10$ g/day
- history of other chronic liver disease (e.g., viral hepatitis, autoimmune hepatitis, primary sclerosing cholangitis, primary biliary cholangitis and overlap syndromes, drug-induced liver injury, hemochromatosis, Wilson's disease, \pm antitrypsin deficiency)
- liver cirrhosis
- any malignancy
- chronic kidney disease
- ... and 5 more (see full listing online)

Locations (2 total)

1st Department of Obstetrics and Gynecology, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece
424 General Military Hospital, Thessaloniki, Greece