

Initial Resuscitation for Acute Kidney Injury in Cirrhosis

NCT06525623

Status	RECRUITING
Phase	Not Applicable
Sponsor	Massachusetts General Hospital
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (5)

- Adult age 18 years or greater
- Signed informed consent form (ICF) by any subject capable of giving consent, or, when the subject is not capable of giving consent, by their legally authorized representatives prior to initiation of any study procedures.
- Admitted to the hospital
- Diagnosis of decompensated cirrhosis (either prior to admission or new diagnosis on admission).
- Presence of acute kidney injury (AKI) as defined by International Club of Ascites (ICA) criteria, defined as SCr increase of e0.3 mg/dL within 48 hours or e50% increase from baseline which is known or presumed to have occurred within the prior 7 days.

Exclusion (14)

- Requiring ≥ 2 liters (L) supplemental oxygen at the time of screening.
- In shock requiring vasopressors (vasoconstrictors for the treatment of AKI such as terlipressin, midodrine, and octreotide are allowed).
- Allergy or other contraindication to IV albumin administration.
- Death, liver transplant, or renal replacement therapy (RRT) expected within 48 hours.
- Patient and/or legally authorized representative unable to provide informed consent.
- ... and 9 more (see full listing online)

Locations (1 total)

Massachusetts General Hospital, Boston, Massachusetts, United States