

Meniscal Infiltration of Corticosteroid Guided With Ultra Sonography

NCT06527235

Status	RECRUITING
Phase	Phase 3
Sponsor	University Hospital, Toulouse
Enrollment	152 participants

Key Eligibility Criteria

Inclusion (6)

- Localized knee pain with tenderness over the medial or lateral joint space reproduced on clinical examination
 - Pain assessed with a VAS score \geq 4/10 despite first-line medical treatment including the use of tier I or II analgesics or NSAIDs.
 - An MRI of the knee performed in the 6 months preceding the operation as part of the pre-treatment assessment
 - Medial or lateral degenerative meniscal lesion on MRI consistent with pain, confirmed by an investigating physician
 - Affiliation to the Social Security
- ... and 1 more (see full listing online)

Exclusion (20)

- Patient under curatorship, guardianship or safeguard of justice
 - Inability to speak, read or write French fluently
 - Patient deprived of liberty
 - Patients with psychiatric pathology
 - Patient who has had an MRI showing an unstable meniscal lesion: complete vertical tear of more than 10 mm in length, capsulomeniscal disinsertion of more than 10 mm in length, complex tear, T2 hypersignal tear of liquid type with passage of liquid testifying to the spreading of the edges, tear with displaced meniscal fragment, lesion of a posterior meniscal brake, lesion of the meniscotibial or menisconfemoral attachment
- ... and 15 more (see full listing online)

Locations (1 total)

UHToulouse, Toulouse, France