

the Study of the Association Between the Gut Microbiota and Central Serous Chorioretinopathy

NCT06527326

Status	RECRUITING
Phase	Not Applicable
Sponsor	Second Affiliated Hospital, School of Medicine, Zhejiang University
Enrollment	400 participants

Key Eligibility Criteria

Inclusion (4)

- Age 18 to 60 years
- Diagnosis of acute central serous chorioretinopathy (CSC) (affected eye: ` FFA and ICGA showing typical CSC manifestations, a presence of subretinal fluid (SRF) at the macular fovea, b disease duration d 6 months)
- Best Corrected Visual Acuity (BCVA): 53 to 86 letters
- Other: No significant systemic diseases, signed informed consent, and able to commit to follow-up

Exclusion (5)

- Presence of significant systemic diseases
- BCVA of 20/200 or worse
- Presence of choroidal or retinal atrophy in the macular fovea of the affected eye
- Presence of choroidal neovascularization (CNV), age-related macular degeneration (AMD), polypoidal choroidal vasculopathy (PCV), diabetic retinopathy (DR), choroidal hemangioma, pigment epithelial detachment (PED), etc., in the affected eye
- High myopia in the affected eye

Locations (1 total)

2 nd Affiliated Hospital, School of Medicine, Zhejiang University, China, Hanzhou, Zhejiang, China