

A Study of Bupivacaine Liposome Injection in the Treatment of Pain After Thoracoscopic Surgery

NCT06529432

Status	RECRUITING
Phase	Phase 2
Sponsor	Jiangsu HengRui Medicine Co., Ltd.
Enrollment	96 participants

Key Eligibility Criteria

Inclusion (6)

- Subjects who are willing to strictly follow the clinical trial protocol to complete this study and voluntarily sign informed consent;
- Elective surgical subjects undergoing lobectomy by single-aperture thoracoscope under general anesthesia;
- age ≥18 years old, Male or female;
- BMI 18-30 kg/m²;
- ASA Physical Status Classification I-II;
- ... and 1 more (see full listing online)

Exclusion (26)

- Participants with any of the following criteria were excluded from the study
- Pre-existing and combined diseases:
 - (1) Subjects with a history of myocardial infarction or unstable angina pectoris, or a history of severe arrhythmias such as atrioventricular block of degree II and above, or NYHA grade II and above in the 6 months before randomization; (2) Subjects with a history of ischemic stroke or transient ischemic attack (TIA); (3) Subjects with psychiatric disorders (such as schizophrenia, depression, etc.) and cognitive dysfunction; (4) Subjects with sensory disorders such as hyperalgesia; (5) Subjects with other physical pain which may affect the evaluation of postoperative pain; (6) Subjects with airway or spinal anatomic factors caused by obstruction of ventilation, bronchiectasis, severe intraoperative thoracic adhesion, etc.
- Laboratory and other tests:
 - Abnormal laboratory results during screening.
- ... and 21 more (see full listing online)

Locations (1 total)

Zhongshan Hospital Affiliated to Fudan University, Shanghai, Shanghai Municipality, China