

Single vs. Dual Antiplatelet Therapy in Elderly or HBR Patients Undergoing Percutaneous Intervention With DCB (PICCOLETO IV-EPIC 38)

NCT06535568

Status	RECRUITING
Phase	Not Applicable
Sponsor	Fondazione Ricerca e Innovazione Cardiovascolare ETS
Enrollment	576 participants

Key Eligibility Criteria

Inclusion (6)

- Male and female patients who meet the following criteria:
- Age \geq 75 years or age \geq 18 years at high bleeding risk;
- Successful PCI with Essential Pro DCB just performed, in 1, 2 or 3 coronary vessels;
- Stable or unstable coronary syndromes;
- De novo coronary lesions in vessels with diameter \geq 2.0 and \leq 4.0 mm (visual estimation);
- ... and 1 more (see full listing online)

Exclusion (26)

- Stent implantation during index or recent ($<$ 6 months) procedure;
- Known (and untreatable) hypersensitivity or contraindication to aspirin, heparin, clopidogrel, paclitaxel or contrast media, or any of their excipient which cannot be adequately pre-medicated;
- Pregnancy at the time of hospitalization;
- Patients participating in another clinical study in which an investigational drug or device was administered within 30 days of screening or within the 5 half-lives of the study drug, whichever is longer;
- ST-elevation myocardial infarction;
- ... and 21 more (see full listing online)

Locations (1 total)

IRCCS Auxologico Milano, Milan, MI, Italy