

Comparison of Palatal Wound Healing in Diabetic and Non-diabetic Patients

NCT06540690

Status	RECRUITING
Phase	Not Applicable
Sponsor	Universidade Estadual Paulista Júlio de Mesquita Filho
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (7)

- Patients with at least 18 years, systemically healthy, with good oral hygiene, assessed by plaque index and gingival index less than 25% (Ainamo, Bay, 1975);
- Patients with no morphological or pathological conditions on the palatine donor area;
- Patients who present indication for extraction and ridge preservation;
- The tooth included in the study, as well as, the adjacent teeth do not present loss of periodontal insertion;
- Patients who agreed to and sign the formal consent to participate in the study after receiving an explanation of risks and benefits from an individual who was not a member of the present study (Resolution no. 118 - May, 2012, and Ethics and Code of Professional Conduct in Dentistry - 118/12).

... and 2 more (see full listing online)

Exclusion (8)

- Patients with systemic problems (cardiovascular, blood dyscrasias, immunodeficiency, and diabetes, among others) that will contraindicate the surgical procedure;
- Patients taking medications known to interfere with the wound healing process or that contraindicate the surgical procedure;
- Smokers patients;
- Pregnant or lactating patients;
- Patients who had had periodontal surgery on the study area;

... and 3 more (see full listing online)

Locations (2 total)

College of Dentistry - São José dos Campos, Sao Paulo State University, São José dos Campos, São Paulo, Brazil
Mauro Pedrine Santamaria and Ana Carolina Ferreira Bonafe, São José dos Campos, São Paulo, Brazil