

# Efficacy of Probucol Combined with Statins Treatment for Ischemic Stroke

NCT06604117

---

**Status** RECRUITING  
**Sponsor** Xuanwu Hospital, Beijing  
**Enrollment** 200 participants

## Key Eligibility Criteria

---

### Inclusion (5)

- Aged 18 years or older.
- Diagnosed with ischemic stroke (IS) confirmed by cranial CT/MRI.
- Onset of stroke within the last 30 days.
- Evidence of atherosclerosis (AS) in at least one artery (carotid, coronary, aorta, renal, or peripheral arteries) identified through neck CTA, coronary CTA, or ultrasound examination of lower extremity arteries.
- Signed informed consent.

### Exclusion (15)

- History of allergy to Probucol or statins.
- Non-atherosclerotic arterial stenosis, such as vasculitis, moyamoya disease, or arterial dissection.
- Potential cardiac embolic sources, such as atrial fibrillation, artificial heart valves, endocarditis, or patent foramen ovale.
- Known bleeding tendencies or hemorrhagic diseases, such as thrombocytopenia (platelet count  $< 100 \times 10^9/L$ ), hemorrhagic stroke, or gastrointestinal bleeding.
- Severe myocardial diseases such as myocardial infarction (MI) or myocarditis.

... and 10 more (see full listing online)

## Locations (1 total)

---

Xuanwu Hospital, Xicheng District, Beijing Municipality, China