

# A Study Evaluating the Safety and Efficacy of the 3M™ V.A.C. Peel and Place Dressing

NCT06625385

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Solventum US LLC
Enrollment	90 participants

## Key Eligibility Criteria

---

### Inclusion (6)

- Subject is at least 22 years old at the time of consent.
  - Subject or legally authorized representative (LAR) is able to provide informed consent.
  - Subject has a wound deemed appropriate for treatment with a V.A.C.® Peel and Place dressing (used in conjunction with 3M™ V.A.C.® Therapy), according to the instructions for use for the dressing, including the following wound types:
    - an open wound: traumatic wound, dehisced surgical wound within 30 days of surgery, burn, venous ulcer, diabetic ulcer, or pressure ulcer Note: Prior to initial placement, the wound may be debrided, as clinically indicated
    - a closed or covered wound secured with sutures or staples: closed surgical incision, skin flap closure, or skin graft (recipient site)
- ... and 1 more (see full listing online)

### Exclusion (30)

- Subject is pregnant or lactating prior to application of the initial dressing. \\*
  - \\*Women who have had surgical sterilization by a medically accepted method (ie, tubal ligation, hysterectomy, or oophorectomy) or are post-menopausal, defined as not having menstruation for \> 12 months will not be required to undergo pregnancy testing.
  - Subject is participating in another interventional clinical study or was enrolled in a clinical trial within the last 30 days before screening.
  - Subject has been diagnosed with a malignancy in the wound.
  - Subject has untreated osteomyelitis or untreated cellulitis in the wound.
- ... and 25 more (see full listing online)

## Locations (3 total)

---

IU Health Methodist Hospital, Indianapolis, Indiana, United States  
Washington University - Barnes Jewish Hospital, St Louis, Missouri, United States  
St. Luke's University Hospital, Bethlehem, Pennsylvania, United States