

Study on the Role of Human Serum Albumin in Large Acute Ischemic Stroke of Anterior Circulation After Thrombectomy

NCT06629116

Status	RECRUITING
Phase	Not Applicable
Sponsor	Tingyu-Yi
Enrollment	100 participants

Key Eligibility Criteria

Inclusion (8)

- Age 18-80 years old.
 - Acute ischemic stroke patients with NIHSS score ≥ 6 .
 - Pre-stroke mRS (modified Rankin Scale) score ≤ 1 .
 - Onset of symptoms to presentation within 24 hours, including wake-up strokes or strokes without witnessed onset; the time of symptom onset is defined as the "last seen normal" time.
 - Confirmed by CTA/MRA/DSA to have anterior circulation large vessel occlusive ischemic stroke (occlusion of the internal carotid artery or M1 or M2 segments of the middle cerebral artery), responsible for acute ischemic stroke signs and symptoms.
- ... and 3 more (see full listing online)

Exclusion (19)

- Intracranial hemorrhage confirmed by head CT or MRI.
 - Pre-stroke mRS score ≥ 2 .
 - Severe allergy or absolute contraindication to iodine-based contrast agents.
 - Systolic blood pressure ≥ 185 mmHg or diastolic blood pressure ≥ 110 mmHg, and inability to control with antihypertensive medication.
 - Blood glucose < 50 mg/dl (2.8 mmol/L) or > 400 mg/dl (22.2 mmol/L) and difficult to correct.
- ... and 14 more (see full listing online)

Locations (1 total)

Zhangzhou Municipal Hospital, Zhangzhou, Fujian, China