

Estimate the Safety and Effectiveness of Adjuvanted Influenza Vaccine Among Asian Elderly People When Compared to Non-adjuvanted Vaccines

NCT06676644

Status	RECRUITING
Phase	Not Applicable
Sponsor	National Taiwan University Hospital
Enrollment	200 participants

Key Eligibility Criteria

Inclusion (2)

- Adults aged 65 years or above
- Presence of at least 1 chronic medical condition associated with increased risk of complicated influenza, including hypertension, chronic obstructive pulmonary disease (group A, B, E), asthma, congestive heart failure (ACC/AHA stage A, B, C, D), coronary arterial disease, diabetes mellitus, chronic liver disease (Child-Pugh score A, B, C) or chronic kidney disease (stage 1~4), neuromuscular disorders, hematological or solid organ malignancies, recipient of hematopoietic stem cell transplantation or solid organ transplantation.

Exclusion (3)

- Contraindications to influenza vaccination including history of severe adverse event after vaccination, such as anaphylaxis or Guillain-Barre Syndrome, or a severe bleeding tendency
- Prior receipt of influenza vaccination during the current flu season.
- Determination by the Principal Investigator (PI) that the participant is ineligible for enrollment based on clinical assessment.

Locations (1 total)

National Taiwan University Hospital, Taipei, Taiwan