

Efficacy of the Entire Papilla Preservation Technique With and Without the Use of L-PRF as a Regenerative Material for the Treatment of Infrabony Defects

NCT06694168

Status	RECRUITING
Phase	Not Applicable
Sponsor	Universidad de los Andes, Chile
Enrollment	30 participants

Key Eligibility Criteria

Inclusion (3)

- Patients diagnosed with periodontal disease (loss of interproximal clinical attachment loss in 2 non-adjacent teeth or clinical attachment loss of ≥ 3 mm associated with a PD > 3 mm in two or more teeth) will be included.
- After re-evaluation (6 weeks after non-surgical periodontal therapy) presented residual probing depths equal to or greater than 6 mm.
- Residual pocket is associated with interproximal intrabony periodontal defects of at least 2 walls, with a depth ≥ 3 mm and an angulation $\geq 40^\circ$ determined on a periapical radiograph.

Exclusion (4)

- Patients with an underlying decompensated pathology such as diabetes and hypertension.
- Patients taking medications that interfere with the healing process such as bisphosphonates.
- Patients with diseases that alter bone metabolism such as osteoporosis.
- Pregnant or breastfeeding women.

Locations (1 total)

CESA (centro de salud de la clínica odontológica de la Universidad de los Andes), Santiago, San Bernardo, Chile