

# Effectiveness of Anti-Inflammatory Drugs for Preventing Pain Versus Treating Pain During Medical Abortion From 13 to 22 Weeks at Hung Vuong Hospital

NCT06696573

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Le Thi Kieu Trang
<b>Enrollment</b>	180 participants

## Key Eligibility Criteria

---

### Inclusion (6)

- Pregnant women aged 18 years or older.
  - No mental illness.
  - Gestational age between 13 and 22 weeks (based on the last menstrual period with a regular menstrual cycle and accurate recall of the last menstrual period, or based on a first-trimester ultrasound) with an indication of stillbirth, fetal malformation requiring termination, or voluntary abortion with counseling and understanding of the abortion procedure.
  - Willing to participate in the study.
  - Capable of reading and signing the informed consent form.
- ... and 1 more (see full listing online)

### Exclusion (9)

- Medical conditions contraindicating medical abortion: bleeding disorders, anticoagulant use, moderate to severe anemia.
  - Cardiovascular conditions: heart failure, mitral valve stenosis, embolism, uncontrolled hypertension.
  - Other medical conditions: renal failure, adrenal gland disorders, severe asthma, liver and biliary diseases, glaucoma.
  - Active pelvic inflammatory disease or acute genital infections.
  - Previous uterine surgery.
- ... and 4 more (see full listing online)

## Locations (1 total)

---

Hung Vuong Hospital, Ho Chi Minh City, Vietnam

---

<https://clinicaltrials.gov/study/NCT06696573>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).