

Best Hypnotic Drug Choice for Rapid Sequence Induction in the Operating Room

NCT06733129

Status	RECRUITING
Phase	Phase 3
Sponsor	Nantes University Hospital
Enrollment	1,218 participants

Key Eligibility Criteria

Inclusion (10)

- age between 18 - 80 years' old
- female* and male
- e risk factor of aspiration of gastric contents defined as
- preoperative fasting period of less than 6 hours,
- occlusive syndrome, functional ileus, vomiting episode within the last 12 hours,
- ... and 5 more (see full listing online)

Exclusion (12)

- predicted impossible tracheal intubation (e 1 of the following criteria: patient with known intubation complications, Mallampati score IV, Thyromental Distance \geq 4.0 cm, Mouth Opening \leq 3 cm, Sternomental Distance \leq 12.5 cm, significant modification of the airway due to congenital, cancer, trauma or burning lesions (non-exhaustive list)) [30]
- preoperative arterial hypotension (MAP \leq 65 mmHg or under catecholamine)
- preoperative respiratory distress syndrome (SpO₂ \leq 90% in room air)
- contraindications to the use of ketamine and/or propofol and/or NMB:
- allergy to the active substance or to one of the excipients or to soy or peanuts,
- ... and 7 more (see full listing online)

Locations (20 total)

Chr Angers, Angers, France
CHRU de Brest Hôpital Cavale Blanche, Brest, France
Ch Louis Pasteur, Chartres, France
... and 17 more locations