

# Intravenous Dexmedetomidine as an Adjuvant to Locoregional Anesthesia in Pediatric Surgery

NCT06736483

---

Status	RECRUITING
Phase	Phase 2
Sponsor	University Hospital, Montpellier
Enrollment	200 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Age 1 to 7 years old male or female
- To be hospitalized for scheduled urovisceral orchidopexy or peritoneovaginal canal surgery (inguinal hernia or hydrocele cure) on an outpatient basis
- Local-regional anesthesia via Transversus Abdominis Plane (TAP) block and/or pudendal block
- National health insurance coverage
- Have obtained signed informed consent from holders of parental authority
- ... and 2 more (see full listing online)

### Exclusion (8)

- Patient under 1 or over 8 years old
- Patients with allergies to local anesthetics
- Patient with a contraindication to locoregional anesthesia: coagulation disorder or infection (fasciocutaneous) in the puncture zone
- Patients with a contraindication to dexmedetomidine (hypersensitivity to the active ingredient or one of the excipients, advanced heart block (level 2 or 3), uncontrolled hypotension, acute cerebrovascular pathologies)
- Patients with delayed psychological development, cognitive or behavioral disorders, or severe neurological pathology.
- ... and 3 more (see full listing online)

## Locations (1 total)

---

UH of Montpellier, Montpellier, France, France