

Performance and Safety of LightForce® Therapy Lasers on Shoulder Soft Tissue Inflammation Pain Reduction

NCT06759935

Status	RECRUITING
Phase	Not Applicable
Sponsor	DJO UK Ltd
Enrollment	130 participants

Key Eligibility Criteria

Inclusion (5)

- Patient male or female with age e18 years old
- Patient with diagnosis of shoulder soft tissue inflammation due to SAIS/RCT (as confirmed by physical examination and/or imaging) to be treated by LightForce® Therapy Lasers according to its indications.
- Patient suffering from shoulder soft tissue inflammation due to SAIS/RCT pain for more than 3 months prior to enrollment
- Pain (either persistent or during activities) score reported by the subject at baseline e 40 mm measured on VAS
- Patient able to provide written informed consent

Exclusion (15)

- Patient with musculoskeletal pathological conditions not to be treated with/contraindication to the use of LightForce® Therapy Lasers according to its intended use and indications
- Patients who are taking drugs that have heat or light sensitive contraindications, such as but not limited to certain types of steroids
- Patients who are administered with corticosteroids, should discontinue the treatment at least 2 weeks prior to study treatment start
- Pregnant females or females of childbearing potentially planning to become pregnant during the study participation
- Patients who had systemic inflammatory conditions (i.e.rheumatoid arthritis, polymyalgia rheumatica)

... and 10 more (see full listing online)

Locations (10 total)

Cabinet Attal, Serris, France
Fisioterapia Carioni, Cassano d'Adda, Milano, Italy
Casertafisio, Caserta, Italy
... and 7 more locations