

# Comparison of Pitavastatin Plus Ezetimibe Versus High-Intensity Statin Therapy on Risk of New-Onset Diabetes Mellitus

NCT06767774

---

Status	RECRUITING
Phase	Phase 4
Sponsor	Korea University Anam Hospital
Enrollment	2,000 participants

## Key Eligibility Criteria

---

### Inclusion (27)

- Age Requirement:
- Patients aged 18 years or older.
- Glycemic Status:
- Patients who are not taking oral hypoglycemic agents (OHAs) and meet all of the following criteria:
- Fasting glucose less than 126 mg/dL.\*

... and 22 more (see full listing online)

### Exclusion (17)

- Patient is pregnant or breastfeeding or of childbearing potential.
- Requires concomitant administration of strong inhibitors of CYP3A4 (itraconazole, ketoconazole, protease inhibitors, erythromycin, clarithromycin, telithromycin, and nefazodone) or CYP2C9 (relative contraindication not dependent on CYP450 statins).
- Chronic kidney disease (eGFR  $< 30$  ml/min/1.73m<sup>2</sup>) or dialysis-dependent renal failure.
- Uncontrolled hypothyroidism.
- Personal or family history of an inherited muscle disorder.

... and 12 more (see full listing online)

## Locations (1 total)

---

Korea University Anam Hospital, Seoul, South Korea