

Intra-Arterial Thrombolysis After SUCCESSful Reperfusion in Anterior Circulation Ischemic Stroke

NCT06768138

Status	RECRUITING
Phase	Phase 3
Sponsor	Central Hospital, Nancy, France
Enrollment	626 participants

Key Eligibility Criteria

Inclusion (9)

- Age \geq 18 years
- Pre-stroke mRS 0-2
- Acute ischemic stroke with anterior circulation large vessel occlusion defined as intracranial internal carotid artery, M1, or M2 occlusion proven on CT or MRI
- NIHSS score \geq 5 at admission
- Acute reperfusion strategy started within 24h after stroke onset according to the international guidelines
- ... and 4 more (see full listing online)

Exclusion (11)

- Person who do not speak French
- Contraindications for intra-arterial thrombolysis: Platelet count $<$ 100 000/mm³, INR $>$ 1.7, AOD use $<$ 48h or biological confirmation of activity and effective heparin treatment
- Bleeding-risk complications during the mechanical thrombectomy procedure (e.g carotid dissection, complicated femoral approach)
- Bleeding-risk complications consecutive to a fall associated with stroke
- More than 5 thrombectomy device
- ... and 6 more (see full listing online)

Locations (1 total)

CHRU Nancy, Nancy, France