

# Assessing Tenapanor as a Treatment of CF-related Constipation.

NCT06810167

---

Status	RECRUITING
Phase	Phase 3
Sponsor	Massachusetts General Hospital
Enrollment	25 participants

## Key Eligibility Criteria

---

### Inclusion (11)

- Having confirmed cystic fibrosis (either by sweat chloride or genetic testing)
  - Meeting criteria for CFrC
  - Must include 2 of the following, with or without abdominal pain for at least 3 months, with symptom onset at least 6-months prior:
    - Straining in at least 25% of defecations
    - Bristol Stool Scale 1-2 more than 25% of defecations (change in stool form)
- ... and 6 more (see full listing online)

### Exclusion (7)

- Use of any antibiotic to treat infection within the 4-weeks prior to study initiation (stable azithromycin dosed 3-times weekly for lung function is to be allowed)
  - Inability to discontinue standing bowel regimen (including fiber, stool softener, as well as either osmotic or stimulant laxative, pro-kinetic serotonergic agents, or other therapy) 2-weeks prior to study drug initiation (with ability to use osmotic laxative therapy as rescue therapy only).
  - Severe CFrC as determined by study team
  - Prior tenapanor usage
  - Hospitalization within 4-weeks prior to study initiation.
- ... and 2 more (see full listing online)

## Locations (1 total)

---

Massachusetts General Hospital, Boston, Massachusetts, United States