

Gene and Molecular Pathways of Ozone Treatment Response in Gynecological Tumor Patients With Chronic Pelvic Pain Secondary to Cancer Treatment

NCT06816095

Status RECRUITING
Sponsor Bernardino Clavo, MD, PhD
Enrollment 40 participants

Key Eligibility Criteria

Inclusion (6)

- Adult women (≥ 18 years old) with gynecological tumors treated with radiotherapy-chemotherapy.
 - Cancer disease is stable or in remission.
 - Life expectancy ≥ 6 months.
 - Patients included in the group of patients with pelvic pain must have a clinical, radiological, endoscopic, or histopathological diagnosis that their pain is not secondary to the oncological process.
 - Patients included in the group of patients with pelvic pain must have pain for ≥ 3 months duration, with an intensity ≥ 3 on the Visual Analog Scale (VAS), or classified as toxicity \geq Grade-2 of the CTCAE v.5.0 of the National Cancer Institute of the USA.
- ... and 1 more (see full listing online)

Exclusion (5)

- Age < 18 years old.
- Severe psychiatric disorders.
- Inability to complete the quality of life questionnaires.
- Active neoplasia requiring recent initiation (< 3 months) of systemic or local treatment.
- Life expectancy (for any reason) < 6 months.

Locations (1 total)

Hospital Universitario de Gran Canaria Dr. Negrín, (FIISC), Las Palmas, Las Palmas, Spain