

# Efficacy and Safety of Cocktail of Ropivacaine for Local Infiltration Analgesia in Patients Undergoing Laparoscopic Cholecystectomy

NCT06819748

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Sichuan Provincial People's Hospital
Enrollment	210 participants

## Key Eligibility Criteria

---

### Inclusion (3)

- Patients undergoing daytime laparoscopic cholecystectomy in Sichuan Provincial People's hospital a Age: 18-70 years
- American society of Anesthesiologists physical status classification system I-II c 18 kg/d BMI d 30 kg/j d No communication barriers, able to understand the research process and the use of pain scale.
- Signed informed consent

### Exclusion (4)

- Drug allergy related to this study
- History of chronic pain, long-term use of analgesic drugs (equivalent to e 10mg oxycodone per day), alcohol abuse, gastrointestinal bleeding or perforation after application of non steroidal anti-inflammatory drugs
- Patients with ischemic heart disease, peripheral arterial vascular or cerebrovascular disease, or patients with pulmonary heart disease, active peptic ulcer or gastrointestinal bleeding, or patients with inflammatory bowel disease
- Patients taking monoamine oxidase inhibitors or within 2 weeks after discontinuation d Preoperative pain score (NRS ) was greater than or equal to 4 points e Participated in other clinical studies within three months

## Locations (1 total)

---

Sichuan Provincial People's Hospital, Chengdu, Sichuan, China