

# COMPARISON BETWEEN EFFICACY OF NOREPINEPHRINE AND PHENYLEPHRINE BOLUSES for PREVENTION OF SPINAL ANAESTHESIA INDUCED HYPOTENSION IN OBSTETRICAL PATIENTS UNDERGOING EMERGENCY CESAREAN SECTION-A Double Blind Randomized Controlled Trial

NCT06836986

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Dow University of Health Sciences
Enrollment	124 participants

## Key Eligibility Criteria

---

### Inclusion (3)

- American society of anesthesiologists II and III patients
- Age 18-40years
- Gestational age 32 weeks and above

### Exclusion (7)

- Patients with hypertensive disorders of pregnancy having baseline systolic blood pressure greater than 160 mmHg and diastolic blood pressure greater than 99 mmHg.
- Baseline mean arterial pressure less than 70 mmHg.
- Antepartum hemorrhage/intraoperative blood loss greater than 1000ml
- history indicative of cardiovascular or neurological disease.
- known fetal abnormality.
- ... and 2 more (see full listing online)

## Locations (1 total)

---

Dr Ruth km pfau civil hospital karachi Dow University of health sciences, department of anesthesiology, Karachi, Sindh, Pakistan