

Clinical Features, Current Treatment and Clinical Outcomes in Patients With INR-CAD: a Cohort Study

NCT06845410

Status	RECRUITING
Sponsor	Peking Union Medical College Hospital
Enrollment	120 participants

Key Eligibility Criteria

Inclusion (5)

- years of age or older, male or female.
- Negative results of urine or blood pregnancy test for females with childbearing potential (not post-menopausal or surgically sterile).
- Meeting the clinical diagnostic criteria for INR-CAD, including: (1) Angiographic evidence of coronary lesions (e 50% diameter stenosis, de novo or restenotic); (2) Evidence of chronic inflammation within 24 months: (A) Positive inflammatory markers (erythrocyte sedimentation rate [ESR], high-sensitivity C-reactive protein [hs-CRP], interleukin-6 [IL-6], tumor necrosis factor-alpha [TNF- α] et al; at least twice, e 12 weeks apart), or (B) Positive autoantibodies (at least twice, e 12 weeks apart), or (C) Established diagnosis of chronic inflammatory diseases (autoimmune disease, systemic vasculitis, psoriasis, tuberculosis, et al), or (D) Receiving immunosuppressive therapy (glucocorticoids, immunosuppressive agents, et al).
- NOT meeting the clinical diagnostic criteria for IR-CAD, including: (1) Hospitalization due to myocardial ischemia, including: (A) Typical symptoms of angina (Canadian Cardiovascular Society [CCS] III-IV), and (B) Non-invasive evidence of myocardial ischemia; (2) Angiographic evidence of new or worsened coronary lesions (de novo or restenotic) considered relevant to myocardial ischemia, which occurred: (A) Within 6 months of last coronary angiography in any patients, or (B) Within 12 months of last coronary angiography in patients receiving immunosuppressive therapy within 24 months.
- Received, or are receiving, or will receive the 24-month clinical follow-up defined by the clinical follow-up protocol for INR-CAD.

Exclusion (10)

- Other moderate to severe heart diseases (congenital heart disease, valvular heart disease, myocarditis, cardiomyopathy, pericardial diseases, pulmonary hypertension, heart failure, arrhythmia, et al).
- Active malignancy (diagnosed within 12 months or with ongoing requirement for treatment).
- Vital organ failure.
- Life expectancy $<$ 1 year.
- In pregnancy or breast-feeding, or with intention to be pregnant during the study period.
- ... and 5 more (see full listing online)

Locations (1 total)

Peking Union Medical College Hospital, Beijing, Beijing Municipality, China

<https://clinicaltrials.gov/study/NCT06845410>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at ClinicalTrials.gov. Generated by ClinicalTrialsFinder.org.