

A Study of 36-Week Refill Exchanges of Port Delivery System (PDS) With Ranibizumab in nAMD

NCT06847542

Status	RECRUITING
Phase	Phase 3
Sponsor	Hoffmann-La Roche
Enrollment	250 participants

Key Eligibility Criteria

Inclusion (6)

- Initial diagnosis of nAMD within 24 months prior to screening
- Previous treatment with at least 3 anti-vascular endothelial growth factor (VEGF) IVT injections for nAMD per standard of care within 6 months prior to screening
- Demonstrated response to prior anti-VEGF IVT treatment since diagnosis
- Availability of historical VA data obtained at or after nAMD diagnosis and prior to the first anti-VEGF treatment for nAMD
- Availability of historical OCT image data obtained at or after nAMD diagnosis and prior to the first anti-VEGF treatment for nAMD
- ... and 1 more (see full listing online)

Exclusion (52)

- A. Prior Ocular Treatment
- Study Eye:
- History of vitrectomy surgery, submacular surgery, or other surgical intervention for age-related macular degeneration (AMD)
- Previous treatment with corticosteroid intravitreal injection
- Previous intraocular device implantation
- ... and 47 more (see full listing online)

Locations (53 total)

Medizinische Universität Innsbruck, Innsbruck, Austria
Kepler Universitätskliniken GmbH - Med Campus III, Linz, Austria
Landeskrankenhaus Mistelbach, Mistelbach, Austria
... and 50 more locations