

Zinc Oxide Ozonated Olive Oil in LSTR

NCT06854744

Status	RECRUITING
Phase	Not Applicable
Sponsor	Tanta University
Enrollment	30 participants

Key Eligibility Criteria

Inclusion (6)

- Bilateral mandibular primary molars showing signs and symptoms of pulp necrosis or chronic pulp infection with/without hemorrhage, suppuration, or purulence.
 - Spontaneous pain or tenderness to percussion. 4- Presence of chronic apical abscess, sinus tract, or grade I mobility. 5-The tooth should be restorable. 6- Radiographic characteristics are:
 - Widening of Periodontal ligament space.
 - Radicular involvement.
 - Furcation involvement less than or equal to half of the shortest root in the vertical dimension
- ... and 1 more (see full listing online)

Exclusion (6)

- Physiologic root resorption is more than one-third of its length.
 - Excessive internal resorption.
 - Non-restorable teeth exhibiting greater than grade I mobility.
 - Perforation into the bifurcation.
 - Any underlying pathological lesion.
- ... and 1 more (see full listing online)

Locations (2 total)

Faculty of dentistry Tanta university, Tanta, Gharbia Governorate, Egypt
Nouran Esmail Mohamed Shalaby, Menouf, Egypt