

# Comparison of Impar Ganglion Block and Sacral Erector Spinae Plane Block in Patients with Coccygodynia

NCT06858917

---

|            |                                       |
|------------|---------------------------------------|
| Status     | RECRUITING                            |
| Phase      | Not Applicable                        |
| Sponsor    | Ba_ak_ehir Çam & Sakura City Hospital |
| Enrollment | 130 participants                      |

## Key Eligibility Criteria

---

### Inclusion (3)

- Those who have had chronic coccygodynia for at least 3 months
- Being between the ages of 18-65
- Patients who do not respond to conservative treatments (analgesics, physical therapy modalities etc.) and are scheduled for interventional pain procedures

### Exclusion (9)

- Patients who have undergone surgery to the coccyx region
- Having undergone an interventional pain procedure in the lumbar, sacral and coccyx regions within the last year
- Patients with systemic infection, local infection at the application site, and coagulopathy
- Pregnant and breastfeeding patients
- Having endocrine or rheumatological disease that may affect the musculoskeletal system in the relevant region and lead to diagnostic complexity (such as spondyloarthritis)
- ... and 4 more (see full listing online)

## Locations (1 total)

---

Ba\_ak\_ehir Çam and Sakura City Hospital, Istanbul, Ba\_ak\_ehir, Turkey (Türkiye)