

Effectiveness of Decision Support for Cardiovascular Risk Management in People With Cardiovascular Disease

NCT06871514

Status	RECRUITING
Phase	Not Applicable
Sponsor	dr.Frank L.J. Visseren
Enrollment	1,200 participants

Key Eligibility Criteria

Inclusion (9)

- Established ASCVD
- At least 30 days since last CVD event and/or the diagnosis of ASCVD. If no CVD events have occurred, imaging confirming ASCVD must have been conducted at least 30 days prior.
- Documented ASCVD (defined according to the 2021 European Society of Cardiology guideline), which includes ASCVD established clinically or demonstrated unequivocally by imaging:
- Clinically documented ASCVD includes previous myocardial infarction, acute coronary syndrome, coronary revascularization (Percutaneous Coronary Intervention (PCI) or Coronary Artery Bypass Surgery (CABG)), and other arterial revascularization procedures, ischemic stroke or transient ischemic attack, and peripheral artery disease (from Fontaine stage II). Angina pectoris (stable) without imaging evidence of atherosclerosis does not qualify as ASCVD.
- ASCVD unambiguously identified through imaging, includes significant stenosis ($\geq 50\%$) on coronary angiography, computed tomography angiography, or carotid ultrasound. It also includes aortic aneurysms measuring ≥ 3 cm. Only Carotid Intima-Media Thickness measurements (cIMT), Coronary Artery Calcium scoring or abnormal ankle-brachial index scores without evidence of stenosis, do not qualify as ASCVD.

... and 4 more (see full listing online)

Exclusion (10)

- Patients currently participating in other interventional medication studies, or studies that directly affect the therapy plan
- Remaining life expectancy of less than 2 years as assessed by a consulting healthcare professional (these patients have no indication for cardiovascular risk management)
- Patients for whom individual risk predictions with the SMART2 model are not feasible:
- Systolic Blood Pressure (SBP) < 90 mmHg or > 200 mmHg
- Total cholesterol < 2.5 mmol/L or > 8 mmol/L

... and 5 more (see full listing online)

Locations (8 total)

Noordwest Ziekenhuisgroep, Alkmaar, Netherlands
Meander Medisch Centrum, Amersfoort, Netherlands
Deventer Ziekenhuis, Deventer, Netherlands

... and 5 more locations

<https://clinicaltrials.gov/study/NCT06871514>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).