

Efficacy of Adding Oral Amisulpride to Dual Prophylaxis for Postoperative Nausea and Vomiting in Patients at High Risk for Nausea and Vomiting Undergoing Gynecological Surgery

NCT06887621

Status	RECRUITING
Phase	Phase 2
Sponsor	Instituto do Cancer do Estado de São Paulo
Enrollment	276 participants

Key Eligibility Criteria

Inclusion (3)

- Laparoscopic hysterectomy to treat benign conditions.
- High risk for PONV according to the Apfel Score: scores 3 or 4.
- American Society of Anesthesiology (ASA) physical status: 1 or 2.

Exclusion (17)

- Cognitive or psychiatric conditions impairing consent or compliance.
- Incapability of using the mobile app MyCapp for data collection.
- History of allergy or sensibility to any medication included in the protocol: amisulpride, dexamethasone, ondansetron, fentanyl, midazolam, bupivacaine, morphine, propofol, rocuronium, sevoflurane, ephedrine, metaraminol, remifentanil, metamizole, ketoprofen, sugammadex, dimenhydrinate, pyridoxine hydrochloride, tramadol, dimethicone.
- Inability to swallow medications.
- Current use of typical or atypical antipsychotic medications.

... and 12 more (see full listing online)

Locations (1 total)

Hospital da Mulher, São Paulo, São Paulo, Brazil