

The Effects of Intravenous Iron on Mobility in Elderly Patients Following Hip Fracture Surgery

NCT06898814

Status	RECRUITING
Phase	Phase 4
Sponsor	Soren Overgaard
Enrollment	210 participants

Key Eligibility Criteria

Inclusion (6)

- years of age or older
- Acute proximal femur fracture surgery
- A hemoglobin measurement ≥ 6.5 mmol/L (10.5 g/dL) on day 1 to 5 postoperatively
- Independent prefracture indoor walking ability, indoor NMS ≥ 2
- Ability to speak and understand Danish

... and 1 more (see full listing online)

Exclusion (11)

- Known allergy to intravenous iron
- Residing permanently at a nursing home
- Hematological conditions with a risk of iron overload e.g. haemochromatosis, hemosiderosis, or where alternative treatments are necessary e.g. haematological malignancies
- Other contraindication to iron treatment, e.g. severe liver cirrhosis and hepatitis
- Severe uncontrolled infection as assessed by the responsible clinician (e.g. bacteraemia or sepsis)

... and 6 more (see full listing online)

Locations (3 total)

Department of Orthopedic Surgery and Traumatology, Bispebjerg, Copenhagen University Hospital, Copenhagen NV, Denmark
Department of Orthopedic Surgery and Traumatology, Herlev, Copenhagen University Hospital, Herlev, Denmark
Department of Orthopaedic Surgery, Odense and Svendborg University Hospital, Odense C, Denmark