

Strategies for Antithrombotic Treatment Following Transcatheter Edge-to-Edge Repair in Patients With an Indication for Oral Anticoagulant

NCT06901466

Status	RECRUITING
Phase	Phase 4
Sponsor	Chinese Academy of Medical Sciences, Fuwai Hospital
Enrollment	880 participants

Key Eligibility Criteria

Inclusion (6)

- Successful TEER procedure, defined as technical success per MVARC criteria.
- Indication for long-term OAC.
- Ability and willingness to comply with the trial protocol.
- Provision of written informed consent.
- Women of childbearing potential must use effective contraception from the time of consent until the final dose of antithrombotic therapy.

... and 1 more (see full listing online)

Exclusion (13)

- Severe renal impairment (creatinine clearance < 15 mL/min or requiring dialysis).
- Ongoing postoperative bleeding (defined as overt bleeding with a ≥ 3.0 g/dL drop in hemoglobin or requiring ≥ 3 units of blood transfusion), or vascular complications following the index TEER procedure.
- Platelet count $< 50 \times 10^9$ /L.
- Need for reoperation due to complications of the index TEER procedure.
- Recent (< 12 month) intracranial or intracerebral hemorrhage.

... and 8 more (see full listing online)

Locations (1 total)

Chinese Academy of Medical Sciences, Fuwai Hospital, Beijing, Beijing Municipality, China