

Effectiveness of Pulpotomy in Symptomatic Permanent Teeth With Deep Caries Lesions

NCT06918639

Status RECRUITING
Sponsor Universidad Rey Juan Carlos
Enrollment 50 participants

Key Eligibility Criteria

Inclusion (8)

- Patients over 18 years of age.
 - Patients in good general health, ASA I or ASA II with no contraindications to receive a TPV procedure under local anaesthesia.
 - Tooth with symptomatology of pulpitis, both reversible and irreversible, to be determined by history of spontaneous pain, exacerbation of pain with heat and cold confirmed by the cold test and radiating pain.
 - Tooth without signs of pulp necrosis and absence of sinus tract and abscess.
 - Tooth in occlusion with antagonist tooth and without discomfort on chewing.
- ... and 3 more (see full listing online)

Exclusion (7)

- Patients with allergies or intolerance to any anaesthetic solution.
 - Pregnant patients or patients who suspect pregnancy due to the impossibility of complementary radiographic tests cannot be carried out.
 - Patients with systemic pathology that prevents the use of a vasoconstrictor or with a compromised immune system that may influence the response to treatment.
 - Tooth with an immature apex.
 - Tooth that is not restorable or not viable for preservation.
- ... and 2 more (see full listing online)

Locations (1 total)

Rey Juan Carlos University, Alcorcón, Madrid, Spain